

Forest Medical Group Annual Pill Check Review

| Thank you for completing this form. We are aiming to avoid the need for you to see a clinician in order to reissue your |
|--|
| pill prescription. If you require the use of a scale or blood pressure machine they can be found in the waiting area – we |
| are unable to issue a repeat prescription without up to date blood pressure and weight measurements. Once completed, |
| please email the form to warrenlane.medicalcentre@nhs.net. If we have any problems with re-issuing your pill we will |
| contact you otherwise we will generate a 12 month supply of the contraceptive pill. Please note, it will take 72 hours to |
| generate your prescription. If you are registered for electronic prescriptions we will send your prescription electronically |
| to your nominated pharmacy. |

| PERSONAL DETAILS | | | | | | | |
|--|---|--------------------------------|--------------------------------------|------------|----------------------|----------------------------|--|
| Full Name | Home Number | | | | | | |
| | | | Work Number | | | | |
| Address and Pos | stcode | | Mobile Numb | er | | | |
| | | | Preferred Number | | | | |
| Date of Birth | | | E-mail Address | | | | |
| Height | Feet/inches | cm | Weight | Stones/Ibs | iones/lbs kg | | |
| Blood Pressure F 2 readings ,5 minute | | nachine in the reception. Take | Blood Pressu | re Reading | 2 | | |
| | | | | | | | |
| | | REVIEW | QUESTION | S | | | |
| Please co | mplete the below que | | /es to any of rther if requir | | ing questi | ons, we may contact you to | |
| Have you had | any problems or con | cerns with the pill? | | | | | |
| Are you aware that the contraceptive pill does <i>NOT</i> protect you from sexually transmitted infections (STIs), so you will need to use a condom as well to protect yourself? | | | | | | | |
| Are you breast feeding? | | | | | If Yes age of child- | | |
| Do you suffer from migraines? | | | | | | | |
| | from high blood press | | | | | | |
| - | • | sis (DVT) or pulmonary | | • | | | |
| pulmonary em | bolism (PE) under the | | n thrombosis | (DVT) or | | | |
| Do you have a | ny blood clotting illne | esses / abnormalities? | | | | | |
| Do you have c | r previously had brea | ist cancer? | | | | | |
| Are you a breast cancer BRCA1/BRCA2 gene carrier? | | | | | | | |
| Have you had a change in bleeding pattern or new bleeding? | | | | | | | |
| Do you smoke | ? | | | | Stop Date | ə (MM/YY) | |
| any new healt | n diagnosed with n conditions since you a prescription? | If Yes ple | ease state - | | | | |
| supplements r the surgery? | any medication or ot prescribed by | | ease state - | | | | |
| Are you aware of the alternatives such as long acting reversible contraceptives (LARCs)? (Please see page 3) | | | | | | | |
| Would you like to book a consultation with a clinician to discuss or arrange fitting of a long acting reversible contraceptive? | | | | | | | |

| It is important that you are aware of: | How the pill works What to do if you miss a pill | If you are not aware or if you require further information please click here or you can discuss further with a clinician | |
|--|--|--|--|
| | That the contraceptive pill may not work if you have diarrhoea or vomiting | I can confirm I understand the information Would you like to discuss this further with a clinician? | |
| | The risks of contraceptive pills | | |

| Name of requested contraceptive pill: | | | | | | |
|--|---------|--|--|--|--|--|
| I can confirm that the information provided is accurate and to the best of my knowledge. I have read and understood the information provided to me regarding contraception and the options in relation to long acting reversible contraception | | | | | | |
| Signature of patient: | Date: | | | | | |
| For office use: Calculated BMI: | Signed: | | | | | |
| Last pill review consultation with clinician (MM/YY): Outcome: | Role: | | | | | |
| | Date: | | | | | |

Г

Please remember your next cervical smear test, book in for an appointment if you are due (EVERY 3 years for women aged between 25-49 and EVERY 5 years for women aged between 50-64) Please click here for further information.

| LARCs→ | | | Intrauterine device | Progesterone |
|--|--|--|---|---|
| | | (IUS) | (IUD) | only injections |
| What is it? | A small, flexible rod inserted under the skin of your upper arm. | A small, T-shaped plastic device is placed into the uterus (womb) | A small plastic and copper device is placed into the uterus (womb) | An injection that slowly releases progestogen |
| How does it work? | It slowly releases the hormone progestogen. It stops ovulation, thickens cervical mucus to stop sperm reaching an egg and thins the lining of the uterus to prevent a fertilised egg implanting | The plastic device releases the hormone progestogen. This thins the lining of the uterus to prevent a fertilised egg implanting and thickens cervical mucus to prevent sperm reaching an egg | The copper prevents sperm from surviving and alters your cervical mucus to prevent sperm reaching an egg. An IUD may also stop a fertilised egg implanting in the uterus | It releases the hormone progestogen which stops ovulation, thickens cervical mucus to prevent sperm reaching an egg and thins the lining of the uterus to prevent a fertilised egg implanting |
| How long does it last for? | 3 years | 5 years. | 5-10 years depending on type. If fitted after age 40, it can stay in place until after the menopause when contraception is no longer needed | Repeat injections every 8-12 weeks depending on type. The injection can't be removed from the body so any side effects may continue for as long as it works and for some afterwards |
| Chances of getting pregnant? | Over 99% effective once fitted. Less than 1 in 100 implant users will get pregnant in a year | Over 99% effective once fitted. Less than 1 in 100 IUS users will get pregnant in a year | Over 99% effective once fitted. Less than 1 in 100 IUD users will get pregnant in a year | Typical use, around 94% effective. 6 in 100 injection users will get pregnant in a year. |
| Could it affect changes of getting pregnant in the future? | Your fertility returns to normal as soon as the implant is removed. | Your fertility returns to normal as soon as the IUS is removed. | Your fertility returns to normal as soon as the IUD is removed. | Your periods and fertility may take up to one year to return after stopping the injection. It may take longer for some people. |
| Affect on periods? | Your periods may stop, be irregular or may last longer. | Your periods usually become lighter, shorter and often less painful. They may stop altogether. | Your periods may become heavier or longer or more painful. | Periods often stop but some women experience irregular or persistent bleeding. |
| Unwanted effects? | Acne, fluid retention, breast tenderness. | Risk of ectopic pregnancy is higher if a woman falls pregnant while using an IUS. Infection. Expulsion – IUS may come out. | Risk of ectopic pregnancy is higher if a woman falls pregnant while using a copper coil. Infection. Expulsion – IUD may come out. | May gain weight (2-3kg over a year) May cause thinning of the bones which is reversible on stopping. |